

ZAVADOVSKIY, A.M., kand. tekhn. nauk

Reciprocal effect of turbine stages on the aerodynamic characteristics
[with summary in English]. Teploenergetika 5 no. 8:8-13 Aug '58.
(MIRA 11:8)

1. Tsentral'nyy kotloturbinnyy institut.
(Steam turbines)

SOV/96-58-8-2/22

AUTHOR: Zavadovskiy, A.M., Candidate of Technical Science
TITLE: The Mutual Influence of Turbine Stages on the Aerodynamic
Characteristics (Vzaimnoye vliyaniye turbinnykh stupeney
na aerodinamicheskiye kharakteristiki)

PERIODICAL: Teploenergetika, 1958, Nr 8, pp 8-13 (USSR)

ABSTRACT: A good deal of experimental work has been done on single turbine stages but interaction between stages has been studied less. In order to find out whether the aerodynamic characteristics of individual stages can be applied when stages work in a group, the Central Boiler Turbine Institute carried out the work on experimental turbines that is described in this article. The tests were made on two groups of stages; in the first the ratio of mean diameter to length was 8.8 - 10, and in the second 3.5 - 6.3. In both cases the blades were twisted. The first set-up consisted of three stages, as shown in Fig 1a. Throughout the investigations the outlet edges of the guide vanes were directed along the radius. The axial distance between stages was 20 mm. Arrangements were made to measure the total and static pressure drops and the direction of flow at different places, so as to be able to evaluate

Card 1/5

The Mutual Influence of Turbine Stages on the Aerodynamic Characteristics

SOV/96-58-8-2/22

the structure of the flow. It was found that, under the test conditions used, later stages have practically no influence on earlier ones and, therefore, tests on the second type of blading were made in a two-stage turbine, a diagram of the flow path of which is given in Fig 1b. The characteristics that were investigated are described, and the criteria of similarity are given. In testing groups of stages, measurements were made of the total power, the power on the wheel rims and the isotropic heat-drop corresponding to the static parameters of the working medium at inlet to and outlet from the stages. An equation is given for the heat-drop corresponding to the power of the second stage when two stages are used; a further expression is given for the efficiency of the second stage of a group. In most respects, the procedure and measurements were as described in earlier work (Teploenergetika Nr 10, 1955). In the first series of tests the characteristics of each of the three stages were first individually studied, then in groups of two and finally all three together. The main test results are given. The changes in the degree

Card 2/5

The Mutual Influence of Turbine Stages on the Aerodynamic Characteristics

SOV/96-58-8-2/22

of reaction at different positions on the blades for a two-stage turbine are plotted in Fig 2. The experimental points on the curves show that the reactions remain practically unaltered whether the stages work separately or together. Graphs of pressure-change on the stage radius appear in Figs 3 and 4, and these too do not depend on whether the stages work singly or together. This also applies to outlet angles of the flow, as will be seen from Fig 5. Thus, with a well-designed flow path, operating near optimum conditions, preceding and succeeding stages make no difference. Curves of change in the total pressure and the outlet angles of flow are given in Fig 6 over the height of stage 2 with various gaps between stage 2 and stage 1. Power losses arise because of leakage through the gaps, as is well shown by Fig 7, which relates the influence of gap length to the change in total pressure over the height of the blades. Impaired flow structure reduces the utilisation of kinetic energy in the stage. Data on the flow of working medium is shown in Figs 8 and 9,

Card
3/5

The Mutual Influence of Turbine Stages on the Aerodynamic Characteristics

SOV/96-58-8-2/22

for various values of radial gap. The flow is much the same whether the stage works individually or in a group. The following practical conclusions are drawn from the work. The aerodynamic characteristics of single stages, as obtained in experimental turbines, can be used when these stages are intermediate ones. The efficiency of the multi-stage flow path depends on the conditions of transition of flow from stage to stage. High leakages past the gaps of the preceding stage impair the flow structure and increase power losses.

Card 4/5

SOV/96-58-8-2/22
The Mutual Influence of Turbine Stages on the Aerodynamic Characteristics

This deterioration increases with the length of the radial gap in the preceding stage; hence these leakages past the gaps are more important in multi-stage than in single-stage turbines.

Card 5/5

There are 9 figures, 2 Soviet references.

ASSOCIATION: Tsentral'nyy kotloturbinnyy institut (Central Boiler Turbine Institute)

- | | |
|---|-----------------------|
| 1. Turbines--Aerodynamic characteristics distribution | 2. Turbines--Pressure |
| 3. Turbines--Test methods | 4. Turbines--Power |

ZAVADOVSKIY, A.M., kand.tekhn.nauk.

Effect of gaps in the joints of segments of rims of runner wheels
on the characteristics of turbine stages. Energomashinostroenie 3
no.8:26-27 Ag '57. (MIRA 10:10)

(Turbines)

ZAVADOVSKIY, A.M., kandidat tekhnicheskikh nauk; BABENKO, Kh.L., inzhener.

Some considerations concerning S.V. Grishchuk's article.
Energomashinostroenie 3 no.9:48 S '57. (MIRA 10:10)
(Gas turbines)

ZAVADOVSKIY, A.M.

ZAVADOVSKIY, A.M., kandidat tekhnicheskikh nauk.

Profile and tip energy losses in a turbine stage [with summary in English]. Teploenergetika 4 no.8:15-18 Ag '57. (MLRA 10:9)

1. TSentral'nyy kotloturbinnyy institut.
(Turbines)

ZAVADOVSKIY, A.M.
BABENKO, Kh.L., inzh.; ZAVADOVSKIY, A.M., kand.tekhn.nauk.

Measures for increasing economic operation of steam turbines.
Elek.sta. 28 no.9:3-6 S '57 (MIRA 10:11)
(Steam turbines)

96-1-7/31

AUTHORS: Zavadovskiy, A.M., Candidate of Technical Sciences and
Babenko, Kh.L., Engineer.

TITLE: The Influence of Leakage on the Operation of a Turbine
Stage (Vliyaniye protechek na rabotu turbinnoy stupeni)

PERIODICAL: Teploenergetika, 1958, Vol.5, No.1, pp. 28 - 31 (USSR)

ABSTRACT: In turbine stages of the type illustrated in Fig. 1, there is an axial gap between the end surface of the diaphragm and the roots of the working blades. The flow through the blading thus has a leakage path into the space surrounding the bucket-wheel. If there are equalising apertures in the bucket-wheel, the working medium in the stage can flow both through the gap and through the apertures. Tests made with an unbladed disc, later confirmed by tests on an experimental turbine, showed that leakage through the peripheral annular gap between the root zone and the diaphragm results from the pumping effect of the rotating disc. Leakage influences the main flow. Gas leaking back from the bucket-wheel space into the flow part alters its direction of motion and is accelerated mainly at the expense of the energy of the main flow. The influence of this kind of leakage would be expected to decrease with increase in the degree of reaction of the stage. Gas leaking from the flow path into the bucket-

Card1/3

The Influence of Leakage on the Operation of a Turbine Stage. 96-1-7/31

wheel space causes hardly any distortion of the flow structure, and moderate leakage scarcely impairs the stage efficiency. Tests were made at the Central Boiler Turbine Institute (TsKTI) to investigate the operation of a stage with the two kinds of leakage. The blading used was that described in a previous article in *Teploenergetika*, 1957, No.6. The gaps considered are those indicated in Fig. 1. The tests were first made on a bucket-wheel without equalising holes and later on a wheel with five equalising holes initially of 15 mm diameter and later of 25 and 35 mm diameter. Graphs of the stage efficiency for various amounts of leakage into the flow path from the space are given in Fig.2. Similar curves for stages with different degrees of reaction are given in Figs. 3 and 4. This kind of leakage affects the stage efficiency differently, depending on the clearance in the blade zone. This is seen in Fig.5, which graphs stage efficiency as a function of leakage for various clearances. The small influence of leakage into the bucket-wheel space from the flow path is confirmed by the additional data graphed in Fig.6, which was obtained for different gap sizes and degrees of reaction. The curve given in Fig. 9 shows that the steam-handling capacity of the stage is little affected by leakage.

Card2/3

The Influence of Leakage on the Operation of a Turbine Stage. 96-1-7/31

There are 9 figures and 2 Slavic references.

ASSOCIATION: TsKTI

AVAILABLE: Library of Congress.

Card 3/3


GUKASOVA, Yekaterina Aleksandrovna; ZHUKOVSKIY, Mikhail Isaakovich;
ZAVADOVSKIY, Anatoliy Mikhaylovich; ZYSINA-MOLOZHEN, Larisa
Mikhaylovna; SKHAR', Nikolay Akimovich; TYRYSHKIN, Vsevolod
Georgiyevich; ZHUKOVSKIY, V.S., prof., doktor tekhn.nauk, red.;
KUTATELADZE, S.S., prof., doktor tekhn.nauk, red.; ZHITNIKOVA,
O.S., tekhn.red.

[Aerodynamic improvement of bladed apparatus of steam and gas
turbines] Aerodinamicheskoe sovershenstvovanie lopatochnykh
apparatov parovykh i gazovykh turbin. Pod red. V.S.Zhukovskogo
i S.S.Kutateladze. Moskva, Gos.energ.izd-vo, 1960. 340 p.

(MIRA 13:7)

(Steam turbines)

(Gas turbines)



The action of synthetic androstosterone and testosterone on the growth of the combs of young roosters. B. M. Zayarnitskii and R. G. Nersisyanova Zayarnitskaya. *Ibid.* *Ibid.*, *in press*. U. R. S. S. R. 3, 25 (1947). Androstosterone in daily doses of more than 0.1 mg. increased the growth of the combs of 40-60 day-old roosters. This growth effect, however, rapidly disappeared and after 4-6 days androstosterone was ineffective. Testosterone was without effect even when 1 mg. was administered daily. It is assumed that testosterone and androstosterone are without effect or have only slight effect on young roosters under a definite age. The physiological effectiveness and the standardization of the male generative-gland hormones. *Ibid.* 29-30.—The "capon-comb units" defined by Huttenand and Ruzicka for testosterone and androstosterone were confirmed. Observation of the behavior of capons after increased hormone doses showed that in addition to the quantitative difference in the action of androstosterone and testosterone there also existed a qualitative difference. Androstosterone regulates, in the first place, the development of the secondary sexual characteristics while the action of testosterone is directed more at the sexual behavior of the animal. When androstosterone was administered immediately after castration, even with a dose as small as 28% a 20% increase in comb growth was obtained. It was also shown that normal pullets can be used for the quantitative estimation of male generative gland hormones. *Through Chem. Zvest.* 1938, 1, 3363 and 3367. M. G. Moore

11F

ca

CHICKENS AS TEST OBJECTS FOR EARLY DIAGNOSIS OF PREGNANCY IN MARES. H. M. ZAVADORSKI and R. G. NEMENOVNA-ZAVADORSKIYA. *Russk. vet. zap.* 1937, 110-18 (1937). — Injections of blood serum of pregnant mares into hens or cocks 30-60 days old and weighing 200-600 g. produced an increase in the comb size even on the second day after the injection. This reaction has been used as a pregnancy test. In rare cases the comb of the chickens did not react to the gonadotropic substance. In these cases a final diagnosis was made on the 6th day after the beginning of the injections when the genital system became markedly more mature than that of the controls. S. A. Corson

ASTM-SLA METALLURGICAL LITERATURE CLASSIFICATION

29

Antagonistic effect of androsterone on hyperthyroid molting of hens. B. M. Zavadovskii and R. G. Nemny-anova-Zavadovskaya. *Bull. Biol. Med. expd. U. R. S. S. S.* 3, 119-20(1937).—A single feeding of dry thyroid gland (10-20 g.) leads to marked and lasting (up to 7 days) molting in hens and castrated roosters but produces very little or no change in the plumage of normal roosters. Injection of male sex hormones into castrated roosters restores their normal resistance to hyperthyroid molting. Injections of synthetic androsterone simultaneously with thyroid feeding to normal hens prevented or markedly decreased molting. The quantity of androsterone injected did not influence the degree of molting but significantly affected the time of onset of molting, the larger doses producing a longer delay. Thyroid feeding did not inhibit the effect of androsterone on comb growth. S. A. C.

ASW-SLA METALLURGICAL LITERATURE CLASSIFICATION

1930-1939

1940-1949

1950-1959

1960-1969

1970-1979

1980-1989

1990-1999

2000-2009

2010-2019

2020-2029

2030-2039

2040-2049

2050-2059

2060-2069

2070-2079

2080-2089

2090-2099

2100-2109

2110-2119

2120-2129

2130-2139

2140-2149

2150-2159

2160-2169

2170-2179

2180-2189

2190-2199

2200-2209

2210-2219

2220-2229

2230-2239

2240-2249

2250-2259

2260-2269

2270-2279

2280-2289

2290-2299

2300-2309

2310-2319

2320-2329

2330-2339

2340-2349

2350-2359

2360-2369

2370-2379

2380-2389

2390-2399

2400-2409

2410-2419

2420-2429

2430-2439

2440-2449

2450-2459

2460-2469

2470-2479

2480-2489

2490-2499

2500-2509

2510-2519

2520-2529

2530-2539

2540-2549

2550-2559

2560-2569

2570-2579

2580-2589

2590-2599

2600-2609

2610-2619

2620-2629

2630-2639

2640-2649

2650-2659

2660-2669

2670-2679

2680-2689

2690-2699

2700-2709

2710-2719

2720-2729

2730-2739

2740-2749

2750-2759

2760-2769

2770-2779

2780-2789

2790-2799

2800-2809

2810-2819

2820-2829

2830-2839

2840-2849

2850-2859

2860-2869

2870-2879

2880-2889

2890-2899

2900-2909

2910-2919

2920-2929

2930-2939

2940-2949

2950-2959

2960-2969

2970-2979

2980-2989

2990-2999

3000-3009

3010-3019

3020-3029

3030-3039

3040-3049

3050-3059

3060-3069

3070-3079

3080-3089

3090-3099

3100-3109

3110-3119

3120-3129

3130-3139

3140-3149

3150-3159

3160-3169

3170-3179

3180-3189

3190-3199

3200-3209

3210-3219

3220-3229

3230-3239

3240-3249

3250-3259

3260-3269

3270-3279

3280-3289

3290-3299

3300-3309

3310-3319

3320-3329

3330-3339

3340-3349

3350-3359

3360-3369

3370-3379

3380-3389

3390-3399

3400-3409

3410-3419

3420-3429

3430-3439

3440-3449

3450-3459

3460-3469

3470-3479

3480-3489

3490-3499

3500-3509

3510-3519

3520-3529

3530-3539

3540-3549

3550-3559

3560-3569

3570-3579

3580-3589

3590-3599

3600-3609

3610-3619

3620-3629

3630-3639

3640-3649

3650-3659

3660-3669

3670-3679

3680-3689

3690-3699

3700-3709

3710-3719

3720-3729

3730-3739

3740-3749

3750-3759

3760-3769

3770-3779

3780-3789

3790-3799

3800-3809

3810-3819

3820-3829

3830-3839

3840-3849

3850-3859

3860-3869

3870-3879

3880-3889

3890-3899

3900-3909

3910-3919

3920-3929

3930-3939

3940-3949

3950-3959

3960-3969

3970-3979

3980-3989

3990-3999

4000-4009

4010-4019

4020-4029

4030-4039

4040-4049

4050-4059

4060-4069

4070-4079

4080-4089

4090-4099

4100-4109

4110-4119

4120-4129

4130-4139

4140-4149

4150-4159

4160-4169

4170-4179

4180-4189

4190-4199

4200-4209

4210-4219

4220-4229

4230-4239

4240-4249

4250-4259

4260-4269

4270-4279

4280-4289

4290-4299

4300-4309

4310-4319

4320-4329

4330-4339

4340-4349

4350-4359

4360-4369

4370-4379

4380-4389

4390-4399

4400-4409

4410-4419

4420-4429

4430-4439

4440-4449

4450-4459

4460-4469

4470-4479

4480-4489

4490-4499

4500-4509

4510-4519

4520-4529

4530-4539

4540-4549

4550-4559

4560-4569

4570-4579

4580-4589

4590-4599

4600-4609

4610-4619

4620-4629

4630-4639

4640-4649

4650-4659

4660-4669

4670-4679

4680-4689

4690-4699

4700-4709

4710-4719

4720-4729

4730-4739

4740-4749

4750-4759

4760-4769

4770-4779

4780-4789

4790-4799

4800-4809

4810-4819

4820-4829

4830-4839

4840-4849

4850-4859

4860-4869

4870-4879

4880-4889

4890-4899

4900-4909

4910-4919

4920-4929

4930-4939

4940-4949

4950-4959

4960-4969

4970-4979

4980-4989

4990-4999

5000-5009

5010-5019

5020-5029

5030-5039

5040-5049

5050-5059

5060-5069

5070-5079

5080-5089

5090-5099

5100-5109

5110-5119

5120-5129

5130-5139

5140-5149

5150-5159

5160-5169

5170-5179

5180-5189

5190-5199

5200-5209

5210-5219

5220-5229

5230-5239

5240-5249

5250-5259

5260-5269

5270-5279

5280-5289

5290-5299

5300-5309

5310-5319

5320-5329

5330-5339

5340-5349

5350-5359

5360-5369

5370-5379

5380-5389

5390-5399

5400-5409

5410-5419

5420-5429

5430-5439

5440-5449

5450-5459

5460-5469

5470-5479

5480-5489

5490-5499

5500-5509

5510-5519

5520-5529

5530-5539

5540-5549

5550-5559

5560-5569

5570-5579

5580-5589

5590-5599

5600-5609

5610-5619

5620-5629

5630-5639

5640-5649

5650-5659

5660-5669

5670-5679

5680-5689

5690-5699

5700-5709

5710-5719

5720-5729

5730-5739

5740-5749

5750-5759

5760-5769

5770-5779

5780-5789

5790-5799

5800-5809

5810-5819

5820-5829

5830-5839

5840-5849

5850-5859

5860-5869

5870-5879

5880-5889

5890-5899

5900-5909

5910-5919

5920-5929

5930-5939

5940-5949

5950-5959

5960-5969

5970-5979

5980-5989

5990-5999

6000-6009

6010-6019

6020-6029

6030-6039

6040-6049

6050-6059

6060-6069

6070-6079

6080-6089

6090-6099

6100-6109

6110-6119

6120-6129

6130-6139

6140-6149

6150-6159

6160-6169

6170-6179

6180-6189

6190-6199

6200-6209

6210-6219

6220-6229

6230-6239

6240-6249

6250-6259

6260-6269

6270-6279

6280-6289

6290-6299

6300-6309

6310-6319

6320-6329

6330-6339

6340-6349

6350-6359

6360-6369

6370-6379

6380-6389

6390-6399

6400-6409

6410-6419

6420-6429

6430-6439

6440-6449

6450-6459

6460-6469

6470-6479

6480-6489

6490-6499

6500-6509

6510-6519

6520-6529

6530-6539

6540-6549

6550-6559

6560-6569

6570-6579

6580-6589

6590-6599

6600-6609

6610-6619

6620-6629

6630-6639

6640-6649

6650-6659

6660-6669

6670-6679

6680-6689

6690-6699

6700-6709

6710-6719

6720-6729

6730-6739

6740-6749

6750-6759

6760-6769

6770-6779

6780-6789

6790-6799

6800-6809

6810-6819

6820-6829

6830-6839

6840-6849

6850-6859

6860-6869

6870-6879

6880-6889

6890-6899

6900-6909

6910-6919

6920-6929

6930-6939

6940-6949

6950-6959

6960-6969

6970-6979

6980-6989

6990-6999

7000-7009

7010-7019

7020-7029

7030-7039

7040-7049

7050-7059

7060-7069

7070-7079

7080-7089

7090-7099

7100-7109

7110-7119

7120-7129

7130-7139

7140-7149

7150-7159

7160-7169

7170-7179

7180-7189

7190-7199

7200-7209

7210-7219

7220-7229

7230-7239

7240-7249

7250-7259

7260-7269

7270-7279

7280-7289

7290-7299

7300-7309

7310-7319

7320-7329

7330-7339

7340-7349

7350-7359

7360-7369

7370-7379

7380-7389

7390-7399

7400-7409

7410-7419

7420-7429

7430-7439

7440-7449

7450-7459

7460-7469

7470-7479

7480-7489

7490-7499

7500-7509

7510-7519

7520-7529

7530-7539

7540-7549

7550-7559

7560-7569

7570-7579

7580-7589

7590-7599

7600-7609

7610-7619

7620-7629

7630-7639

7640-7649

7650-7659

7660-7669

7670-7679

7680-7689

7690-7699

7700-7709

7710-7719

7720-7729

7730-7739

7740-7749

7750-7759

7760-7769

7770-7779

7780-7789

7790-7799

7800-7809

7810-7819

7820-7829

7830-7839

7840-7849

7850-7859

7860-7869

7870-7879

7880-7889

7890-7899

7900-7909

7910-7919

7920-7929

7930-7939

7940-7949

7950-7959

7960-7969

7970-7979

7980-7989

7990-7999

8000-8009

8010-8019

8020-8029

8030-8039

8040-8049

8050-8059

8060-8069

8070-8079

8080-8089

8090-8099

8100-8109

8110-8119

8120-8129

8130-8139

8140-8149

8150-8159

8160-8169

8170-8179

8180-8189

8190-8199

8200-8209

8210-8219

8220-8229

8230-8239

8240-8249

8250-8259

8260-8269

8270-8279

8280-8289

8290-8299

8300-8309

8310-8319

8320-8329

8330-8339

8340-8349

8350-8359

8360-8369

8370-8379

8380-8389

8390-8399

8400-8409

8410-8419

8420-8429

8430-8439

8440-8449

8450-8459

8460-8469

8470-8479

8480-8489

8490-8499

8500-8509

8510-8519

8520-8529

8530-8539

8540-8549

8550-8559

8560-8569

8570-8579

8580-8589

8590-8599

8600-8609

8610-8619

8620-8629

8630-8639

8640-8649

8650-8659

8660-8669

8670-8679

8680-8689

8690-8699

8700-8709

8710-8719

8720-8729

8730-8739

8740-8749

8750-8759

8760-8769

8770-8779

8780-8789

8790-8799

8800-8809

8810-8819

8820-8829

8830-8839

8840-8849

8850-8859

8860-8869

8870-8879

8880-8889

8890-8899

8900-8909

8910-8919

8920-8929

8930-8939

8940-8949

8950-8959

8960-8969

8970-8979

8980-8989

8990-8999

9000-9009

9010-9019

9020-9029

9030-9039

9040-9049

9050-9059

9060-9069

9070-9079

9080-9089

9090-9099

9100-9109

9110-9119

9120-9129

9130-9139

9140-9149

9150-9159

9160-9169

9170-9179

9180-9189

9190-9199

9200-9209

9210-9219

9220-9229

9230-9239

9240-9249

9250-9259

9260-9269

9270-9279

9280-9289

9290-9299

9300-9309

9310-9319

9320-9329

9330-9339

9340-9349

9350-9359

9360-9369

9370-9379

9380-9389

9390-9399

9400-9409

9410-9419

9420-9429

9430-9439

9440-9449

9450-9459

9460-9469

9470-9479

9480-9489

9490-9499

9500-9509

9510-9519

9520-9529

9530-9539

9540-9549

9550-9559

9560-9569

9570-9579

9580-9589

9590-9599

9600-9609

9610-9619

9620-9629

9630-9639

9640-9649

9650-9659

9660-9669

9670-9679

9680-9689

9690-9699

9700-9709

9710-9719

9720-9729

9730-9739

9740-9749

9750-9759

9760-9769

9770-9779

9780-9789

9790-9799

9800-9809

9810-9819

9820-9829

9830-9839

9840-9849

9850-9859

9860-9869

9870-9879

9880-9889

9890-9899

9900-9909

PROCEDURES AND PROPERTIES INDEX																									
1st and 2nd columns													3rd and 4th columns												
<p><i>ca</i></p> <p>The influence of prolactin on the sexual activity of mares. B. M. Zavadovskii and M. B. Goldberg. <i>Bull. Acad. Sci. USSR Div. Vet. Med. Biol. Sci.</i> 1938, II, 3411. --Previously sterile mares were treated with relatively small doses of prolactin (500-1000 mouse units) was divided into 3 doses administered on the first 3 days of estrus. In 60-65% of cases pregnancy followed. Nymphomania should be treated with larger doses (2000-7000 mouse units). After repeated matings, some of the treated animals became pregnant at once, some the following year. No control expts. in order to det. the percentage of spontaneous recovery from the condition are treated.</p> <p style="text-align: right;">M. G. Moore</p>																									
<p>ASH-SEA METALLURGICAL LITERATURE CLASSIFICATION</p> <p>8304 634100</p> <p>8304 634100</p> <p>8304 634100</p>																									

1ST AND 2ND COLUMNS										PROCESSING AND PROPERTY INDEX										3RD AND 4TH COLUMNS									
COMMON ELEMENT										COMMON TRANSITION METALS										<p>The effect of hormones of the sterin group on the oviduct of the pullet. By M. Zayulovskii, E. G. Nemyayanova, Zayulovskaya and B. P. Roden. <i>Bull. biol. med. exp.</i> U. R. S. S. 640-9(1937); <i>Chem. Zentr.</i> 1938, II, 3104; cf. C. A. 32, 7022. In expts. on hens the injection of solutions of folliculin, folliculin benzoate, androsterone, testosterone or luteosterone produced hypertrophy of the oviduct and growth of the comb. The serum of pregnant mares had a still more pronounced effect than these hormones. Since the different substances of the sterin group produced very similar effects, a common transformation in the ovaries of the animals does not appear to be out of the question.</p> <p style="text-align: right;">M. G. Moore</p>									
A 18-51A METALLURGICAL LITERATURE CLASSIFICATION										E 1111111111111111																			
SIGNATURE										SIGNATURE																			
100000 1111111111111111										100000 1111111111111111																			
100000 1111111111111111										100000 1111111111111111																			

ZAVADOVSKIY, B.M.

TUSHNOV, Mikhail Pavlovich, 1879-1935.

(The problem of spermatoxin and lysates; collection) Pod red. B.M. Zavadovskogo i K.
R. Viktorova. Moskva, Sel'-khozgiz, 1938. 414 p.

The antagonistic effect of androsterone on hyperthyroid
 moult in chicks. H. M. Zavodovskii and E. G. Nersisyanova-Zavodovskaya. *J. THYROID* (U. S. S. R.) 24, 162-
 61 (in English, 191) (1938). The injection of 0.2-0.8 mg.
 of androsterone (I) into normal young hens along
 with the ingestion of a dried thyroid gland prepn. (II) re-
 sults in an inhibition of the hyperthyroid moulting reac-
 tion. I does not inhibit the effect of II on the combs of
 normal hens. S. A. Karjala

11 F

2a

The effect of sex hormones on the quantity of erythrocytes and hemoglobin in the blood of fowl. B. M. Zavadskii and E. P. Rozen (K. A. Timiryazev Biol. Museum, Moscow). *Izvest. Akad. Nauk SSSR, Ser. Biol. Med.* 13, No. 3/4, 30-30 (1942). Sexual dimorphism is seen in the number of erythrocytes in fowl blood, normal cocks having 37%, more than capons or pullets. To det. the nature of this, young cocks, pullets, and adult birds were injected with sex hormones. Injection of folliculin resulted in great variation in blood counts of different birds or the same bird at different times, without regularity. Injection of androstosterone or testosterone in pullets led to marked uniform increase in erythrocytes and hemoglobin, to the level of that in normal cocks; with young capons there was a temporary increase in these effects without lasting effect as in pullets. Injection of gonadostimulatory stallion serum or human placental blood had no effect on adult fowl, but markedly increased erythrocytes and hemoglobin in the blood of pullets or cockerels. The latter showed premature development of seminal vesicles, combs, and sperms. The effects on erythrocytes and hemoglobin are regarded as indirect, acting through influence on the sex glands, and not directly on the blood.

K. Starr Chester

ASH-ILA METALLURGICAL LITERATURE CLASSIFICATION

SECTION DIVISION

COLLECTION

RESEARCH DIVISION

101 AND 210 SERIES										100 AND 210 SERIES									
PROCESSING AND PROPERTY INDEX																			
<div style="position: absolute; top: 10px; left: 10px; font-size: 2em; font-weight: bold;">A4</div> <div style="position: absolute; top: 10px; left: 100px; font-size: 1.5em;">ZAYADOVSKIY, G. M.</div> <div style="position: absolute; top: 300px; left: 300px; font-size: 0.8em;"> <p>Special content of section in relation to reproductive function and content of pregnancy hormones in horses' blood. In G. M. Zayadovskiy and R. M. Neumajskoye-Zayadovskiy: (Soviet Acad. Sci. U.S.S.R., 1960, vol. 227-229). -- The low rate of pregnancy among covered mares in a stud farm in the Czech region, and deficiency of serum-gonadotropin in pregnant mares and the continued presence after resorption of the foetus are discussed and attributed to a deficiency of vitamins A and E in the fodder of the region.</p> </div>																			
AIR-55A METALLURGICAL LITERATURE CLASSIFICATION										6-177072-20 17									
100000 MAP DIV 500										100000 MAP DIV 500									
100000 MAP DIV 500										100000 MAP DIV 500									

BC ZAVODOVSKIY, B.M. A-4

Experiments conducted in vitro indicated that the follicle-stimulating hormone in pregnant-mare serum, B. M. Zavodovski (Comm. Acad. Sci. U.R.S.S., 1943, 47, 213-214), 2-30 young cows treated with group I pregnant-mare serum were identical in wt. and general health with those from 20 controls but differed from them in producing an average of one more offspring per farrow. Data obtained confirm the theory that the optimum effects of the pregnant-mare serum should contain a definite ratio of two gonad-stimulating substances: Factor A, the follicle-stimulating substance, and factor B, which stimulates the ovulation of mature follicles and the formation of corpora lutea. Pregnant-mare serum-II contains the optimum ratio of factors for inducing oestras and serum-I the optimum for stimulating the generative functions of the ovary. A. G. P.

ASB-56A METALLURGICAL LITERATURE CLASSIFICATION

SIGN DIVISION										SIGN ROMANY									
SIGN DIVISION										SIGN ROMANY									
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20

ZAVADOVSKIY, Boris Mikhaylovich; BOBYLEV, P.G., redaktor; VESKOVA, Ye.I.,
tekhnicheskii redaktor.

[Animal and plant; brief introduction to the science of life]
Zhivotnoe i rastenie; malen'koe vvedenie v nauku o zhizni. Izd.
2-oe, dop. i perer. N.G. Mesmanovoi-Zavadovskoi. Moskva, Gos. izd-
vo sel'khoz.lit-ry, 1956. 69 p. (MLRA 9:5)

1, Deystvitel'nyy chlen Vsesoyuznoy akademii sel'skokhozyaystvennykh
nauk imeni V.I. Lenina (for Zavadovskiy).
(Biology)

ZAVADOVSKIY, B.M., akademik; SKVORTSOV, I.M., red.

[The origin of domestic animals] Proiskhozhdenie domashnikh
shivotnykh. Moskva, Gos.izd-vo sel'khoz.lit-ry, 1945. 54 p.
(Domestic animals) (Evolution) (MIRA 13:1)

ZAVADSKIY, B., inzh.

Speeding up the motor-bus transportation. Zhil.-kom.khoz. 9
no.6:29 '59. (MIRA 12:10)
(United States--Traffic regulations)

ZAVADOVSKIY, Boris Mikhaylovich; NESMEYANOVA-ZAVADOVSKAYA, Ye.G.; BOBYLEV,
P.G., redaktor; ZURILINA, Z.P., tekhnicheskiy redaktor

[Origin of domestic animals] Proiskhozhdenie domashnikh zhivotnykh.
Izd. 4-oe, dop. i perer. E.G. Nesmeianovoi-Zavadovskoi. Moskva,
Gos. izd-vo sel'khoz. lit-ry, 1956. 111 p. (MIRA 10:4)
(Paleontology) (Domestic animals)

ZAVADOVSKIY, D. K.

Zavadovskiy, D. K. "X-ray examination of the diaphragm following an operation of diaphragmocrurotomy", Sbornik trudov, posvyashch. prof. Savinykh, Tomsk, 1948, p. 234-39.

So: U-3261, 10 April 1953 (Letopis 'Zhurnal 'nykh Statey, No. 12, 1949).

ZAVADOVSKIY, D. K.

Zavadovskiy, D. K. -- "X-ray observations on stomach patients after balneological treatment in Shira spa, " Sbornik trudov (Tomskiy obl. nauch.-issled. in-t fiz. metodov lecheniya i kurortologii), Vol. VI, 1949, p. 83-88

SO: u-5241, 17 December 1953, (Letopis 'zhurnal Snykh Statey, No. 26, 1949).

ZAVADOVSKIY, D.K.

Some problems in dosimetry of betatron radiation. Med. rad.
no. 5:87-89 '61. (MIRA 14:11)

1. Iz kafedry rentgeno-radiologii i kafedry akusherstva i gineko-
logii Tomskogo meditsinskogo instituta.
(RADIATION---DOSAGE) (BETA RAYS---THERAPEUTIC USE)

ZAVADOVSKIY, D.K.

On the critical remarks on my article, "Some problems in dosimetry
of betatron radiations." Med.rad. 7 no.7:78-81 JI '62.

(MIRA 15:11)

(RADIATION—DOSAGE)

ACC NR: APTC05336

SOURCE CODE: UR/0181/67/009/001/0139/0144

AUTHOR: Zavadskiy, E. A.; Fekidov, I. G.

ORG: Institute of Physics of Metals, AN SSSR, Sverdlovsk (Institut fiziki metallov AN SSSR)

TITLE: Magnetic properties of the alloy FeRh in strong magnetic fields

SOURCE: Fizika tverdogo tela, v. 9, no. 1, 1967, 139-144

TOPIC TAGS: iron base alloy, rhodium containing alloy, saturation magnetization, temperature dependence, critical point, antiferromagnetism, phase transition

ABSTRACT: In view of discrepancies between the results of other investigators, the authors measured in detail the magnetization of FeRh over a wide range of magnetic fields and temperatures. The iron was alloyed with 53 at.% of rhodium in a high-frequency furnace and in an inert atmosphere. The measurements were made on solid samples by an induction method and on powders by means of a pulsed magnetic balance. The two measurement procedures were described by the authors earlier (FIZM v. 12, 832, 1961 and v. 21, 693, 1966). The measurements were made at temperatures 77 - 400K and in magnetic fields up to 330 kOe. The results showed that saturation set in at temperatures above the critical value at which the FeRh goes over from the antiferromagnetic into the ferromagnetic state. The dependence of the critical field on the temperature is a straight line with constant slope in the entire range of temperatures and magnetic fields. Temperature hysteresis of the electric resistivity and of the

Card 1/2

ACC NR: AP7605336

magnetization were clearly observed in the tests at the transition from the antiferromagnetic into the ferromagnetic state. It is pointed out that this linear dependence changes if a change takes place in the magnetic structure at some value of the magnetic field, causing the slope of the line to change. The authors thank A. Ya. Afanas'yev for preparing the alloy and I. I. Kuntsevich for help with the measurements. Orig. art. has: 6 figures and 4 formulas.

SUB CODE: 20/ SUBM DATE: 03Jun66/ ORIG REF: 004/ OTH REF: 009

Card 2/2

BA

A.III - 18

Effect of ascorbic, riboflavin C, and nicotinic acid on development of D-hyperkeratosis. R. N. Matsko and E. V. Zakharenko. *J. Physiol., USSR*, 1951, 27, 233-239. -Rats placed on a diet containing excess amounts of irradiated ergosterol developed signs of D-hyperkeratosis. This could be prevented or cured by administration of carotene but not by ascorbic or nicotinic acid. D. 16, 16477.

ZAVADOVSKAYA, Ye.K.; IVANKINA, M.S.; MELIK-GAYKAZYAN, I.Ya.

Pore formation during annealing of mixed KCl-KBr crystals. Kristallografiia 5 no.2:324-325 Mr-Apr '60. (MIRA 13:9)

1. Tomskiy politekhnicheskii institut.
(Potassium chloride) (Potassium bromide)

ZAVADSKAYA, I.G.; FEL'DMAN, N.L.; KAMENTSEVA, I.Ye.

Carbohydrate content and cold resistance in the cells of higher plants. Dokl. AN SSSR 157 no.4:995-997 Ag '64 (MIRA 17:8)

1. Botanicheskiy institut im. V.L. Komarova AN SSSR i Institut tsitologii AN SSSR. Predstavleno akad-mikon N.M. Sisakyanom.

ZAVADSKAYA, I.O.

Effect of unfavorable soil moisture conditions on the protoplasm
of the epidermis cells in "Viner" barley leaves. Uch. zap. Ped.
inst. Gerts. 178:181-183 '59. (MIRA 14:7)
(Barley) (Plants, Effect of soil moisture on) (Protoplasm)

SKAZKIN, F.D.; ZAVADSKAYA, I.G.

Microsporogenesis in barley as affected by inadequate soil moisture
and nitrogen nutrition. Dokl. AN SSSR 117 no.1:150-152 H-D '57.
(MIRA 11:3)

1. Leningradskiy gosudarstvennyy pedagogicheskiy institut im.
A.I.Gertsena. Predstavleno akademikom A.L.Kursanovym.
(Barley) (Spores (Botany))

ZAVADSKAYA, I.G., Cand Biol Sci -- (diss) "Effect of nitrogen ^{up} -
on certain physiological processes and the formation of
[in barley] pollen ^{under moderate} ~~when there is insufficient~~ [supply of water] ^{to it} ~~to it~~
the soil ^{during} ~~in~~ various periods of development." Len, 1959,
19 pp (Min of Education RSFSR. Len State Pedagogical Inst
in A.I. Gertsen) 150 copies (KL, 35-59, 113)

GOL'TSMAN, Lyubov' Naumovna; kand.ekonom.nauk; ZAVADSKAYA, Irina
Yevseyevna, kand.ekonom.nauk; ORLOVA, Raisa Il'ichna,
nauchnyy sotrudnik; YAMPOL'SKAYA, Tat'yana Georgiyevna,
kand.tekhn.nauk; KHOLMOGOROVA, T.A., red.izd-vo; SHLIKHT,
A.A., tekhn.red.

[Maintaining city streets] Voprosy ekspluatatsii gorodskikh
dorog. Moskva, Izd-vo M-va kommun.khoz.RSFSR, 1959. 88 p.
(MIRA 12:11)

(Streets--Maintenance and repair)

ZAVADSKAYA, M. S.

"Trachoma in Belorussia and Means of Eradicating It." Cand Med Sci,
Minsk State Medical Inst, 2 Dec 54. (SB, 18 Nov 54)

Survey of Scientific and Technical Dissertations Defended at USSR Higher
Educational Institutions (11)

SO: Sum. No. 521, 2 Jun 55

ZAVADSKAYA, T. I.

PA 31/49T34

USSR/Medicine - Undulant Fever
Medicine - Vaccine Therapy

Nov 48

"The Problem of Using Vaccine for Treating Brucellosis,
T. I. Zavadskaya, Hosp Therapeutics Clinic, Rostov
Med Inst, 7 $\frac{1}{2}$ pp

"Klin Med" Vol XXVI, No 11

Data is based on 60 cases. Concludes that vaccine
cannot be regarded as specific remedy for brucellosis.
It does, however, lower temperature and decrease pain.

31/49T34

ZAVADSKAYA, T.I., dotsent; DOMBROVSKAYA, Ye.A.

Clinical aspects and pathomorphology of acute tuberculous
sepsis. Sov. med. 27 no.2:9-14 F '64.

(MIRA 17:10)

1. Kafedra gospital'noy terapii (zav. - prof. N.M.Ivanov) i
kafedra patologicheskoy anatomii (zav. - prof. I.I. Dorokhov)
Rostovskogo-na-Donu meditsinskogo instituta.

ZAVADSKAYA, T.I., kand.med.nauk

The problem of treatment of diabetic coma; Sov.med. 22 no.6:107-109
Je '58 (MIRA 11:9)

1. Iz kafedry gosital'noy terapii (nav. - prof. N.M. Ivanov)
Rostovskogo meditsinskogo instituta.
(DIABETES MELITUS, compl.
coma, ther. (Rus))

ZAVADSKAYA, T.I. (Rostov-na-Donu); VORONOV, A.S., professor, zaveduyushchiy.

Clinical aspects of thrombosis of the splenic vein. Klin.med. 31 no.7:52-
56 J1 '53. (MLBA 6:9)

1. Kafedra gosspital'noy terapii Rostovskogo meditsinskogo instituta.
(Thrombosis) (Veina--Diseases)

ZAVADSKAYA, T.I., kandidat meditsinskikh nauk (Rostov-na-Donu)

Two cases of cutaneous xanthomatosis. Klin.med. 34 no.4:57-59
Ap '56. (MLRA 10:1)

1. Iz kafedry gosptal'noy terapii (zav. - prof. A.S.Voronov)
Rostovskogo meditsinskogo instituta.

(LIPOIODOSIS,

cutaneous, case reports (Rus))

(SKIN, diseases,

xanthomatosis, case reports (Rus))

ZAVADSKAYA, T. N.

USSR/Geophysics - Physics of the Earth

FD-1720

Card 1/1 : Pub. 45-8/12

Authors : Berdichevskiy, M. N., and Zavadsкая, T. N.

Title : On the formation of an electric field in the earth

Periodical : Izv. AN SSSR, Ser. geofiz., 178-180, Mar-Apr 1955

Abstract : When a constant current is introduced into a circuit consisting of connecting wires, electrodes and the earth, transitional processes take place in the earth resulting in the formation of an electromagnetic field. Having examined the records obtained at a low level of interference from the field of telluric currents, the authors present diagrams in which are depicted eight basic forms of the formation of the elastic field. The diagrams show impulse, calibrating impulse and current impulse.

Institution : Scientific Research Institute of Geophysical Methods of Prospecting, Ministry of Petroleum Industry

Submitted : November 23, 1953

ZAVADSKAYA, T.N.

On transformation of sounding curves. Prikl. geofiz. no. 19:47-56
'58.

(Prospecting--Geophysical methods)

(MIRA 11:4)

ZAVADSKAYA, T.N.

Results of testing the method of average apparent resistance.
Razved. geofiz, no.4:52-58 '65. (MIRA 18:9)

CHERNOMORDIK, A.B.; BASS, T.M.; BASS, M.A.; KOVALENKO, F.N.; ZAVADSKAYA, TS.Ye.

Neomycin-resistant forms of colienterites in children and their treatment. Antibiotiki 10 no.9:859-861 S '65.

(MIRA 18:9)

1. Otdel antibiotikov Kiyevskogo instituta epidemiologii i mikrobiologii.

LAMKINA, V.Yu.; NEMCHINOV, G.A.; ZAVADSKAYA, V.A.

Use of pituitrin in cases of difficult egg-laying. Ptitsse-
vodstvo 8 no.8:45 Ag '58. (MIRA 11:10)

1. Kafedra akusherstva i ginekologii Buryat-Mongol'skogo zoovet-
instituta.

(Eggs--Production)

(Pituitrin)

ZAVADSKIY, A.M.

Craspedacusta marginata (Modser) polyps in Tashkent, Dokl. AN
SSSR 60 no.5:921-923 My '48. (MLRA 10:8)

1. Predstavleno akademikom L.S. Bergom.
(Coelenterata)

ZAVADSKIY, A. M.

PA 68784

USSR/Medicine - Marine Organisms
Medicine - Taxonomy

May 1948

"Craspedacusta Marginata (Modser) Polype in Tashkent,"
A. M. Zavadekiy, 3 pp

"Dok Ak Nauk SSSR" Vol LX, No 5

Subject polyp is only form of hydrozoa possessing
metagenesis. Describes characteristics of rare polyps
found by author in an aquarium in 1941. Gives state
of polype in aquarium and methods employed to study
them. Submitted by Academician L. S. Berg 12 Mar
1948.

FDB

68784

ZAVADSKIY, B., inshener.

Gyrobuz. Zhil.-kom.khoz. 5 no.8:27 '55.
(Switzerland--Motorbuses)

(MLBA 9:3)

ZAVADSKIY, B., inzh.

Antiskid chains. Avt.transp. 38 no.1:58-60 Ja '60.
(MIRA 13:5)
(Automobiles--Apparatus and supplies)

ZAVADSKIY, B.

Diagnosis, Radioscopic

Use of X-rays in medicine and the role of the nurse in roentgenological examination of patients. Med. sestra, No. 4, 1952.

Monthly List of Russian Accessions, Library of Congress, November 1952. UNCLASSIFIED.

ZAVADSKIY, B.

NURSES AND NURSING

Use of X-rays in medicine and the role of the nurse in roentgenological examination of patients. Med. sestra no. 4, 1952.

Monthly List of Russian Accessions, Library of Congress. November, 1952. Unclassified.

ZAVADSKIY, B.F.

GAVRILOV, A.N., doktor tekhnicheskikh nauk, redaktor; POLIAKOV, K.A., professor, retsenzent; ZAVADSKIY, B.F., inzhener, retsenzent; RUSEVICH, I.M., inzhener, redaktor; MODIL', B.I., tekhnicheskii redaktor; TIKHONOV, A.Ya., tekhnicheskii redaktor

[Progressive practice in instrument making] Progressivnaia tekhnologiya priborostroeniia. Moskva, Gos. nauchno tekhn. izd-vo mashinostroita. i sudostroita. lit-ry. No.2. [Perfecting instrument production techniques] Sovershenstvovanie tekhnologii proizvodstva priborov. Pod red. A.N.Gavrilova. 1953. 337 p. (MLRA 8:3)

1. Vsesoyuznoye nauchnoye inzhenerno-tekhnicheskoye obshchestvo mashinostroiteley i priborostroiteley.
(Instruments)

ZAVADSKIY, Boris Iosifovich

[Five years overseas; notes on Canada] Piat' let za okeanom; kanadskie zapiski. Moskva, Sovetskii pisatel', 1961. 349 p.
(MIRA 14:8)

(Canada--Description and travel)

ZAVADSKIY, B. I., ENG.; IL'IN, V. A., ENG.

Steam Boilers

Block mount for PK-10 boiler, Elek, sta., 23, no. 6, 1952.

Monthly List of Russian Accessions, Library of Congress, October 1952. UNCLASSIFIED.

ZAVADSKIY, F. I.; IL'IN, V. A., ENG.

Steam Boilers - Testing

Hydraulic testing of boilers with the help of compressed air. Elek. sta.
23 no. 8, 1952.

Monthly List of Russian Accessions. Library of Congress, November 1952. UNCLASSIFIED.

ZAVADSKIY, B. I.; IL'IN, V. A., Engs.

Steam Boilers - Testing

Hydraulic testing of boilers with the help of compressed air. Elek. sta. 23 no. 8, 1952.

Monthly List of Russian Accessions, Library of Congress, November 1952. Unclassified.

ZAVADSKIY, B. I., ENG.; IL'IN, V. A., ENG.

Steam Boilers

Block mount for PK-10 boiler, Elek, sta., 23. no. 6, 1952

Monthly List of Russian Accessions, Library of Congress October 1952. Unclassified.

ZAVADSKIY, B.I., inzhener; IL'IN, V.A., inzhener.

Installing a boiler on previously prepared post foundation. Elek.sta. 24
no.9:51-52 S '53. (MLRA 6:8)

(Boilers)

1. ZAVADSKIY, B.I., ENG.
2. USSR (600)
4. Steam Boilers
7. Tightening the flange connections of steam cooler of a high-pressure boiler. Elek. sta. 23 no.9, 1952.
9. Monthly List of Russian Accessions, Library of Congress, January 1953. Unclassified.

ZAVADSEIY, B.I.

Automobile trailer. Avtomobilist 1:124-129 '61. (MIRA 15:1)
(Automobiles--Trailers)

ZAVADSKIY, B.S.

Readers discuss "Meditsinskaia Sestra" at a readers' conference in
Kolomna. Med. sestra 19 no.9:41-42 8 '60. (MIRA 13:9)
(NURSES AND NURSING--PERIODICALS)

ZAVADSKIY, B.S. (Kolonna)

"School children at the health station". Med.sestra 15 no.6:29
Je '56. (MIRA 9:8)
(SCHOOL HYGIENE) (BORUTSKA, I.M.) (RED CROSS)

ZAVADSKIY, B.S.

Application of roentgen rays in medicine and role of a nurse
in x-ray examination of patients. Med. sestra, Moskva no. 8:22-26
Aug 1952. (GML 23:1)

1. Roentgen Technician. 2. Kolomna, Moscow Oblast.

ZAVADSKIY, B.S. (Kolonna)

"The road to health, strength, and long life." I.M.Sarkizov-Serazini. Reviewed by B.S.Zavadskii. Fel'd. 1 akush. no.10:62-63
O '55. (MLRA8:12)

(PUBLIC HEALTH) (SARKIZOV-SERAZINI, I.M.)

ZAVADSKIY, B.S.(Kolomna, Moskovskaya oblast')

Character qualifications required of a nurse. Med.sestra no.8:
18-23 Ag '55. (MLRA 8:11)

(NURSING PROFESSION

in Russia, communist moral features)

(ETHICS, MEDICAL

, in Russia, communist moral features of nurse)

ZAVADSKIY, B.S., fel'dsher (Kolonna)

The nurse as promotor of health education. Med. sestra no.10:
28-30 0 '54.

(MLRA 7:12)

(BIOGRAPHIES

Serikova, Klaudiia Grigor'evna)

ZAVADSKIY, E. S.

ZAVADSKIY, B.S.

Brief news. Med. sestra no.8:29-30 Ag '54.

(MLRA 7:8)

1. Predsedatel' soveta meditsinskikh sester.
(NURSES AND NURSING--PERIODICALS)

ZAVADSKIY, B.S. (Kolomna)

Letters to the editor. Med.sestra 18 no.10:44-45 0 '59.

(MIRA 13:1)

1. Predsedatel' Soveta srednikh meditsinskikh rabotnikov.
(KOLOMNA (MOSCOW PROVINCE)--MEDICAL PERSONNEL)

ZAVADSKIY, B.S.

A good tradition. Med. sestra 20 no.9:61-63 S '61. (MIRA 14:10)
(NURSES AND NURSING--PERIODICALS)

AUTHORS: Fakidov, I. G. and Zavaḱskiy, E. A. SOV/126-6-3-28/32

TITLE: Generation of Super-intensive Magnetic Field Pulses
(Polucheniye sverkhsil'nykh impul'snykh magnitnykh
poley)

PERIODICAL: Fizika Metallov i Metallovedeniye, 1958, Vol 6, Nr 3,
p 569 (USSR)

ABSTRACT: In 1929, Academician P. L. Kapitza managed to obtain magnetic surge fields with potentials of up to 3 600 000 Oe and utilised them for studying the galvanometric properties of a large number of metals and some semi-conductors. Earlier, the same author produced a field with a potential of the order of 500 000 Oe in a coil of 1 mm dia. by discharging a powerful battery (Ref 1). The interest in phenomena relating to the influence of strong magnetic fields on the physical properties of metals and semi-conductors has considerably increased since that time. Developments in theoretical physics in recent years led to the conclusion that investigation of the magnetic and galvanometric properties of solid bodies in the field of intensive and super-intensive magnetic fields can yield important information on the shape and topology of energy surfaces of conductivity

Card 1/4

SOV/126-6-3-28/32

Generation of Super-intensive Magnetic Field Pulses

electrons (Ref 2). In the laboratory of electric phenomena of the Institute of Metal Physics, Ural Branch, Ac.Sc. USSR a test rig is at present in operation for generating strong magnetic fields using short current pulses obtained by discharging a condenser battery of 1600 μ F capacity charged to a potential of 3000 V. The discharge of the condenser battery through a coil is periodic with a frequency of 2800 to 3000 c.p.s. and a damping decrement $\Delta = 3$ to 5.5, depending on the number of turns of the coil. This set-up permits generating inside a single-layer coil a magnetic field with a potential of over 500 000 Oe with a degree of uniformity of up to 1.5% inside a cylinder of 6.5 mm dia. and a height of 5 mm (it is mentioned in a footnote that the authors have succeeded in raising this potential up to 700 000 Oe). In Fig.1 an oscillogram is reproduced which shows the dependence on time of the potential of the magnetic field. The authors measured the dependence

Card 2/4

SOV/126-6-3-28/32

Generation of Super-intensive Magnetic Field Pulses

of the electric conductivity of n and p-type germanium at a high frequency ($\rho = 54 \text{ Ohm}\cdot\text{cm}$, $\rho = 58 \text{ Ohm}\cdot\text{cm}$) on the transverse magnetic field for $T = 300, 77$ and 20°K . It was established that $\Delta R_1/R$ of n-type germanium ($\rho = 54 \text{ Ohm}\cdot\text{cm}$) at $T = 20^\circ\text{K}$ is subjected to fluctuations. The results of these measurements and a detailed description of the set-up for measuring intensive magnetic fields will be published in later work. Acknowledgments are made to N. V. Volkenshteyn for supplying the liquid hydrogen and to K. I. Davidenko for carrying out the measurements. There are 1 figure and 6 references, 1 of which is Soviet, 5 English.

(Note: This is a complete translation)

Card 3/4

SOV/126-6-3-28/32

Generation of Super-intensive Magnetic Field Pulses

ASSOCIATION: Institut fiziki metallov Ural'skogo filiala AN SSSR
(Institute of Metal Physics, Ural Branch of the
Ac. Sc., USSR)

SUBMITTED: January 8, 1958

- | | |
|-----------------------------------|----------------------------------|
| 1. Magnetic fields--Development | 2. Magnetic fields--Applications |
| 3. Magnetic fields--Effectiveness | 4. Magnetic fields--Measurement |
| 5. Pulses--Properties | |

Card 4/4

AUTHORS: Fakidov, I. G., ~~Zavadskiy~~, E. A. 56-34-4-56/60

TITLE: Oscillations of the Electric Resistance of n-Type Germanium in Strong Pulse-Like Magnetic Fields (Ostsillyatsiya elektricheskogo soprotivleniya germaniya n-tipa v sil'nykh iapul'snykh magnitnykh polyakh)

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1958, Vol. 34, Nr 4, pp. 1036 - 1037 (USSR)

ABSTRACT: The authors investigated the change of the electric resistance of 3 monocrystalline germanium samples of the n-type in a transversal pulse-like magnetic field with an intensity up to 120 000 Gauss at temperatures of 300,77 and 200K. The magnetic field was produced by means of the discharge of a condenser bank by a solenoid, and in the opening of that solenoid a Dewar flask containing the sample was put up. The germanium samples were of different degrees of purity. In magnetic fields of 25 000 - 120 000 Gauss and at $T = 300\text{K}$, $\Delta R/R_0$ depends linearly on the field intensity in the case of all 3 samples, the 3 angles of gradient of the line are given. At 77°K and in the same interval of the field intensities the linear dependence holds only for 2

Card 1/3

Oscillations of the Electric Resistance of n-Type
Germanium in Strong Pulse-Like Magnetic Fields

56-34-4-56/60

samples. In the third sample that dependence has a curved character with a tendency to saturation. The change of the $\Delta R/R_0$ of the sample number 1 (specific resistance $\rho = 54 \Omega \text{ cm}$) was also investigated at 20°K in the case of field intensities up to 110 000 Gauss. It is interesting that in such a case the resistance of the sample decreases instead of increasing as usual. But in the case of a reduction of the amplitude to zero the resistance of the sample returns to its original value. Besides this fact in the case of such germanium samples of the n-type an oscillation of the electric resistance in the interval of the electric field intensities of 25 000 - 110 000 Gauss was observed. The period of that oscillation is 0,10 Kilogauss and its maximum amplitude $H = 55 \text{ 000 Gauss}$. The author points to different previous works, dealing with the same subject. Data on details of the experiments and on the devices for the production of strong magnetic fields will be published in a later paper. There are 1 figure and 5 references, 1 of which is Soviet.

Card 2/3

Oscillations of the Electric Resistance of n-Type
Germanium in Strong Pulse-Like Magnetic Fields

56-34-4-56/60

ASSOCIATION: Institut fiziki metallov Ural'skogo filiala AN SSSR
(Institute of Physics of Metals, Urals Branch, AS USSR)

1. Germanium crystals--Oscillations

Card 3/3

SOV/126-7-4-24/26

AUTHORS: Fakidov, I.G. and Zavadskiy, E.A.

TITLE: An Induction Method of Measuring the Hall Effect in Strong Pulsed Magnetic Fields

PERIODICAL: Fizika metallov i metallovedeniye, 1959, Vol. 7, Nr 4, pp 637-638 (USSR)

ABSTRACT: In the classical method of measuring the Hall effect a primary current from an external source is passed through a sample in a magnetic field. If the magnetic field is not constant, currents induced in the sample may be used instead of the primary current. A method using varying magnetic fields to measure the Hall constant was first described by Busch et al (Ref 1); they used currents induced on switching on or off of a d.c. electromagnet. The present authors describe an application of the Busch method to strong periodic pulsed magnetic fields and materials of high resistivity such as semiconductors. A sample in the form of a disc of radius r_0 was placed in a coil at right-angles to magnetic force lines (Fig 1). The varying magnetic field induced currents in the disc. For samples of high-resistivity material in the form of thin discs, the surface effects and the

Card 1/3

SOV/126-7-4-24/26

An Induction Method of Measuring the Hall Effect in Strong Pulsed Magnetic Fields

demagnetizing action of induced currents may be neglected. Since the applied magnetic field is of damped oscillatory nature the magnetic induction is given by

$$B = B_m e^{-bt} \sin \omega t \quad (4)$$

where $b = \delta/T$, δ is the logarithmic decrement and T is the time period. The value of the Hall emf between the centre of the disc and its periphery, at the moment of the first maximum $B_{m1} = B_m \exp(-\delta/4)$ of the magnetic induction, is given by

$$V_{x1} = -AR\sigma \omega B_{m1}^2, \text{ where } A = \frac{1}{8} e^{\delta/4} (1 - \tan \varphi),$$

R is the Hall constant, $\tan \varphi = b/\omega$, ω is the angular frequency, σ is the electrical conductivity of the sample. The relationship obtained here was checked on a germanium disc of 11 mm diameter, 1 mm thickness in magnetic fields up to 120 kilogauss and $\omega = 16000 \text{ sec}^{-1}$. The calculated values of the Hall constant R were compared with the results of measurements on a plate of

Card 2/3

SOV/126-7-4-24/26

An Induction Method of Measuring the Hall Effect in Strong Pulsed Magnetic Fields

the same material; the two sets of results agreed satisfactorily. The method can be used also for undamped alternating magnetic fields. Then $b = 0$ and $\varphi = 0$ and the Hall emf between the centre of the disc and its periphery is given by

$$V_x = -\frac{1}{8} R \omega^2 B_m^2 \sin 2\omega t \quad (9)$$

In low-frequency magnetic fields and for thin discs the relationships obtained by the authors are also valid for metals. There is 1 figure and 1 Swiss reference.

ASSOCIATION: Institut fiziki metallov AN SSSR
(Metal Physics Institute, AS USSR)

SUBMITTED: December 4, 1958

Card 3/3

67687

SOV/126-8-4-9/22

24.7600

AUTHORS: Fakidov, I.G., and Zavadskiy, E.A.

TITLE: A Generator of Ultrahigh Pulsed Magnetic Fields

PERIODICAL: Fizika metallov i metallovedeniye, 1959, Vol 8, Nr 4, pp 562-568 (USSR)

ABSTRACT: The pulsed magnetic fields are obtained by discharging a bank of capacitors through special coils. Magnetic fields up to 700 000 oersted can be obtained in this way. A photograph of the apparatus is shown in Fig 1. Sixteen type IM-3/100 capacitors are employed so that 7200 joules can be stored at a nominal voltage of 3 kV. By reducing the resistance and the inductance of the discharge circuit it was possible to increase the percentage of energy used to produce the magnetic field to 17%. The coils can take currents up to 60 000 amp. A block diagram of the apparatus is shown in Fig 2. The bank of capacitors is charged from the high-voltage rectifier through the current limiting resistor R. The discharge takes place through the spherical discharger 3. If necessary, the circuit can be controlled automatically to produce the required current pulses. A drawing of one of the coils is given in Fig 4, and the

Card
1/2

67687

SOV/126-8-4-9/22

A Generator of Ultrahigh Pulsed Magnetic Fields

corresponding magnetic field distribution for a coil with an internal diameter of 16 mm is shown in Fig 5. The magnetic field was measured with a search coil in conjunction with an RC integrator, and the calculated value of 700 000 oersted was confirmed experimentally. The ultrahigh pulsed magnetic fields are being used by the present authors in a study of galvanomagnetic phenomena and of the photogalvanomagnetic effect in various semiconductors. Magnetisation studies on ferrites are also being carried out. A similar apparatus has been built in the low-temperature laboratory of the Moscow State University Professor A.I. Shal'nikov, Corr. Memb.

AS USSR. Acknowledgement is made to I.I. Kuntsevich and A.A. Teterin.

There are 6 figures, 2 tables and 7 references, of which 1 is French, 1 is Soviet and 5 are English.

ASSOCIATION: Institut fiziki metallov AN SSSR
(Institute of Physics of Metals, Ac.Sc. USSR)

SUBMITTED: February 28, 1959

Card 2/2

ZAVADSKIY, E.A.; FAKIDOV, I.G.

Electric conductivity of n-type germanium in strong magnetic fields.
Fiz. met. i metalloved. 10 no.3:495-496 S '60. (MIRA 13:10)

1. Institut fiziki metallov AN SSSR.
(Germanium—Electric properties) (Magnetic fields)

ZAVADSKIY, E. A., Cand. Phys-Math. Sci. (diss) "Galvano-magnetic Properties of Germanium in Strong Impulse Magnetic Fields."

Sverdlovsk, 1961, 17 pp. (Acad. of Sci. USSR, Institute of Physics of Metals) 170 copies (KL Supp 12-61, 251).

89947

9.4300 (1043, 1143, 1150)

S/126/61/011/001/015/019
E052/E314

AUTHORS: Zavadskiy, E.A. and Fakidov, I.G.

TITLE: Electrical Conductivity of n-Ge in Strong Pulsed
Magnetic Fields

PERIODICAL: Fizika metallov i metallovedeniye, 1961,
Vol. 11, No. 1, pp. 145 - 147

TEXT: In a previous paper the authors showed that the relative change in the resistance of n-Ge reaches a saturation value in a strong longitudinal magnetic field, and beginning at a certain value of the field, commences to increase linearly. These results are in agreement with the theoretical predictions of Tsidil'kovskiy and Shirokovskiy (Ref. 2) and Gold and Roth (Ref. 3). The present work is concerned with the variation in the resistivity in a strong transverse magnetic field. n-Ge monocrystalline specimens having resistivities of 2, 30 and 46 ohm.cm at room temperature were used. The specimen dimensions were 9 x 1.5 x 0.8 mm. Current leads covering end surfaces and voltage probes 3.5 mm apart

Card 1/8

89947

S/126/61/011/001/015/019
E032/E314

Electrical Conductivity of n-Ge in Strong Pulsed Magnetic Fields were used. The contacts were ohmic and the measurements were carried out as in Ref. 1. Fig. 1 shows the relative change in the resistivity for the specimen with $\rho = 2 \text{ ohm.cm}$ at different temperatures as a function of the magnetic field (in kOe). The results refer to the case $H \parallel [111]$. In this specimen the mobility of current carriers was $\mu = 3500 \text{ cm}^2/\text{Vsec}$ so that at $H = 200 \text{ kOe} \approx 7.0$. Saturation would be expected to set in at $\mu H \approx 1$. The quantity $(\Delta\rho/\rho_0)_{\text{sat}}$ for $H \parallel [111]$ can be calculated from the formula:

Card 2/8

89947

S/126/61/011/001/015/019
E032/E314

Electrical Conductivity of n-Ge in Strong Pulsed Magnetic Fields

$$\left(\frac{\Delta e_{\perp}}{\rho_0} \right)_{\text{sat}} = \frac{(2\gamma + 1)(4\gamma + 5)}{3\gamma(\gamma + 8)} - 1$$

given in Ref. 2, where $\gamma = m_l/m_t$ and m_l and m_t are the longitudinal and transverse effective masses. Assuming that $\gamma = 17$, one finds that $(\Delta e_{\perp}/\rho_0)_{\text{sat}} = 1.0$. However, complete saturation is not observed. Continuous approach to saturation is, in fact, replaced by a linear increase. A "discontinuity" in the curve is nevertheless observed at $(\Delta e_{\perp}/\rho_0) = 1.2$, i.e. the discontinuity occurs close to the theoretical value of $(\Delta e_{\perp}/\rho_0)_{\text{sat}}$. The "discontinuity" can apparently be explained by the quantisation of the energy of the current carriers in the magnetic field which becomes appreciable for $\hbar w > kT$ where w is the cyclotron frequency and \hbar is the Planck constant divided by 2π . Assuming that

Card 3/8

89947

S/126/61/011/001/015/019
E032/E314

X

Electrical Conductivity of n-Ge in Strong Pulsed Magnetic Fields
the effective mass $m^* = 0.08 m_0$, one finds that $hw = kT$
will occur for $H = 35$ kOe and $T = 58^\circ \text{K}$, while at $T = 77^\circ \text{K}$
the magnetic field should be 45 kOe. It is clear from Fig. 1
that the "discontinuity" does in fact occur at magnetic fields
close to those for which $hw = kT$. According to Argyres
(Ref. 4), the linear relation between $(\Delta \rho_{\perp} / \rho_0)$ and H should
occur in non-degenerate semiconductors when $hw > kT$ and
scattering on phonons terminates. For specimens having
resistivities of 2 ohm.cm or greater, scattering on impurity
ions can dominate only below 20°K . Thus the present results
are in good agreement with those reported in Ref. 4. Fig. 2
shows the plot of $(\Delta \rho_{\perp} / \rho_0)$ as a function of $H(\text{kOe})$ for the
specimen with $\rho = 30$ ohm.cm. In this figure $H \parallel [110]$.
A sharp transition to the linear law is found to occur at
 $T = 58$ and 77°K when $(\Delta \rho_{\perp} / \rho_0) = 2.9$, which is close
to the calculated value $(\Delta \rho_{\perp} / \rho_0) = 3.3$. The latter value
Card 4/8

89947

S/126/61/011/001/015/019

E032/E314

Electrical Conductivity of n-Ge in Strong Pulsed Magnetic Fields was obtained for $\gamma = 17$ from the formula given by Gold and Roth in Ref. 3. The "discontinuity" disappears at high temperatures in weak magnetic fields (up to 18 kOe). The results obtained by the present authors at $T = 94^\circ \text{K}$ are in good agreement with those reported by Herring et al (Ref. 5). The resistivity of some specimens was also measured at $T = 20^\circ \text{K}$. In the case of specimens with $\rho = 30 \text{ ohm.cm}$ with $H \parallel [111]$ the curves of $(\Delta\rho_\perp/\rho_0)$ versus H exhibit the following behaviour: 1) for fields up to 15 kOe the curves approach saturation at $(\Delta\rho_\perp/\rho_0) = 3.0$; 2) between 35 kOe and 110 kOe the curve is linear and $(\Delta\rho_\perp/\rho_0)$ increases from 4 to 20; 3) above 110 kOe the curve becomes nonlinear and the values of $(\Delta\rho_\perp/\rho_0)$ are 25, 29.5 and 31.5 at 140, 170 and 200 kOe, respectively. The Hall effect has also been measured and the results will be reported in the next paper of the present journal.

Card 5/8

89947

S/126/61/011/001/015/019
E032/E314

Electrical Conductivity of n-Ge in Strong Pulsed Magnetic Fields

There are 2 figures and 6 references: 3 Soviet and
3 non-Soviet.

ASSOCIATION: Institut fiziki metallov AN SSSR (Institute of
Physics of Metals of the AS USSR)


SUBMITTED: July 20, 1960

Card 6/8

S/126/61/011/001/615/019
EO32/E314

Electrical Conductivity of n-Ge in Strong Pulsed Magnetic Fields

Fig.1.




Card 7/8

89947

S/126/61/011/001/015/019
E032/E314

Electrical Conductivity of n-Ge in Strong Pulsed Magnetic Fields

Fig.2.



Card 8/8